

Original Article

Comparative study on the efficacy of statin therapy in lowering cholesterol and assessing the risk of metabolic syndrome in patients under statins therapy



Kodipelly Ramana Raju¹, Afreen Sharifa¹, Paspula Soumya^{1,*}, Rumana Khanam¹, Koyala konda Banda Sanjay Bhargav¹



Article info

Received: 09 Jan 2022

Revised: 14 May 2022

Accepted: 01 Jul 2022

Use your device to scan and read the article online



Keywords:

Metabolic Syndrome, Lipid Profile, Statin Therapy, Hypertension

ABSTRACT

The objective of the study is to compare and evaluate the efficacy of atorvastatin (group – A) versus rosuvastatin (group – B) on baseline parameters like lipid profile tests and to assess the risk of metabolic syndrome using a Mets calculator. A total of 100 patients were enclosed in the present study who met the inclusion criteria. They were divided into two groups based on their treatment plan Group A includes 24 males and 26 females while Group B includes 23 males and 27 females. The mean differences before treatment for group A and group B are as follows, HDL (31.52 ± 0.35 and 28.34 ± 0.480), LDL (161.4 ± 1.09 and 163.16 ± 0.94), Total cholesterol (252.82 ± 1.09 and 255.56 ± 1.26) and Triglycerides (214.2 ± 0.86 and 215.98 ± 0.62), VLDL (35.98 ± 0.56 and 36.12 ± 0.43). The mean differences after treatment for group A and group B are as follows HDL (39.92 ± 0.46 and 42.04 ± 0.30), LDL (144.96 ± 0.68 and 138.34 ± 0.73), Total cholesterol (181.48 ± 1.98 vs 174.32 ± 2.08), Triglycerides (185.94 ± 1.22 vs 181.74 ± 1.77), VLDL (27.14 ± 0.21 and 24.72 ± 0.27). Group B ($P=0.001$) exhibited a significantly greater reduction in cholesterol levels as compared to Group A ($P = 0.002$). The reductions in LDL, VLDL, Total Cholesterol, and Triglycerides along with increased HDL levels were found to be significantly more in the Rosuvastatin group. In this study, we observed that patients on Rosuvastatin exhibited better control over lipid profile when compared to patients who are on Atorvastatin. Since, this study was conducted on a smaller number of patients, to make consecutive remarks about the superiority of either of the treatment regimen; further analysis of clinical trials is required for appropriate selection of the best statin therapy.

1. Introduction

Metabolic syndrome is a complex constellation of disorders that increase the risk of coronary heart disease (CHD) [1-3]. Reaven first described the metabolic syndrome in 1988 as "Syndrome X." He noted that risk factors such as dyslipidemia, hypertension, and hyperglycemia clustered together and increased CVD risk [4]. In 1998,

the World Health Organization (WHO) characterized the metabolic syndrome as insulin resistance requiring either a fasting insulin level in the upper quartile in nondiabetic patients [5], a fasting plasma glucose greater than 110 mg/dL, or a plasma glucose greater than 200 mg/dL 2 hours after an oral glucose tolerance test and at least two of the following criteria: 1) hypertension

¹Smt. Sarojini Ramulamma College of Pharmacy, Seshadrinagar, Mahabubnagar, Telangana 509001, India

*Corresponding Author: Paspula Soumya (paspulasoumya07@gmail.com)

requiring systolic or diastolic blood pressures of 160 mm Hg or greater or 90 mm Hg or greater or treatment for hypertension [6]; 2) dyslipidemia defined as triglycerides (TG) greater than 150 mg/dL or HDL cholesterol less than 35 mg/dL in men or less than 39 mg/dL in women, or treatment for dyslipidemia [7]; 3) body mass index (BMI) greater than 30 kg/m² and/or waist-to-hip ratio greater than 0.9 in men, greater than 0.85 in women, or a waist girth of 37 inches or greater in men and 32 inches or greater in women; or 4) microalbuminuria (urinary albumin excretion rate \geq 20 μ g/min or albumin: creatinine ratio \geq 20 μ g/g) [8, 9].

2. Materials and methods

2.1. Study sites, design, and period

The study is a prospective, parallel, open-label observational study conducted in the general medicine department of a 300-bedded multi-specialty tertiary care teaching hospital, for 6 months. A total of 100 prescriptions were included in the study. The study protocol and written informed consent were approved by the institutional ethical committee of the hospital. To consider both inpatients and outpatients of the general medicine ward of SVS hospital.

2.2. Inclusion criteria

- ✓ Patients of any gender of age above 20yrs age were diagnosed with dyslipidemia.
- ✓ Patients with DM, HTN, or any cardiovascular and cerebrovascular disease.
- ✓ Patients who are willing to give their consent for the study.
- ✓ The patient was prescribed any one of the drugs (atorvastatin and rosuvastatin)

2.3. Exclusion criteria

- ✓ Patients who were not treated with statin therapy.
- ✓ All patients with immunodeficient diseases like HIV.

- ✓ Patients with concomitant diseases like T.B.
- ✓ Patients who are not willing to give their consent for the study.
- ✓ Pregnant and lactating women are excluded.

2.4. Data collection

- ✓ Data collection form.
- ✓ Metabolic syndrome calculator.

2.5. Study procedure

- ✓ This is a prospective study where patient data are collected and written in the data collection form. This form contains the demographic details of the patient, chief complaint, past medical, and medication history, family history, laboratory investigations and medication chart.
- ✓ The efficacy study of statins will be compared by taking lipid profile data from patients. The risk of metabolic syndrome is assessed by the use of a metabolic syndrome calculator.
- ✓ The study will be conducted on patients of department general medicine and cardiology where anyone of a statin drug (atorvastatin, rosuvastatin) was prescribed at SVS Medical College hospital. All information relevant to the study using suitable methods for statistical analysis and results will be concluded.

3. Results

3.1. Distribution of patents according to age

During 6 months period, a total of 100 participants were collected. Collected patients were divided into two groups based on their treatment plan as group A (Atorvastatin) and group B (rosuvastatin). More participants were observed in the age group above 60yrs. It has been shown in Table 1 and Figures 1-4. Table 2 shows that before starting the treatment, there was no significant difference between the two groups. Tables 3 and 4 show

the risk assessment of metabolic syndrome in statin therapy.

Table 1. Distribution of Patients according to age

Age category	GroupA (No of Patients)	Group B (No of Patients)
20 - 35	3(6%)	5(10%)
36 - 45	4(8%)	5(10%)
46 - 60	17(34%)	15(30%)
>60	26(52%)	25(50%)
Total	50(100%)	50(100%)




Fig. 1. Distribution of Patients according to age




Fig. 2. Comparison of changes in biochemical parameters for group B

Table 2. Comparison of biochemical parameters

Parameters	Group A	Group B
HDL	31.52±0.35	28.34±0.48
LDL	161.4±1.09	163.16±0.94
Total cholesterol	252.82±1.10	255.56±1.26
Triglycerides	214.2±0.86	215.98±0.62
VLDL	35.98±0.56	36.12±0.43
Cholesterol HDL ratio	8.03±0.09	8.61±0.10

Table 2. The significant difference before and after treatment in group B

Parameters	Before treatment	After treatment	P-value
HDL	28.34±0.48	42.04±0.30	0.0001
LDL	163.16±0.94	138.34±0.73	0.01
Total cholesterol	255.56±1.26	174.32±2.08	<0.0001
Triglycerides	215.98±0.62	181.74±1.77	<0.0001
VLDL	36.12±0.43	24.72±0.27	0.0001
Cholesterol HDL ratio	8.61±0.10	4.15±0.05	<0.0001

Table 3. Significant changes in parameters in statin therapy

Parameters	Category	Atorvastatin	Rosuvastatin
BMI	Above 30 kg/m ²	33	26
	below 30kg/m ²	17	24
Triglycerides	below 150mg/dl	22	15
	above 150mg/dl	28	35
HDL	above 50mg/dl	1	5
	40-49mg/dl	23	27
Blood pressure	below 40mg/dl	26	18
	below 130/85mmHg	15	12
Type II diabetes	above 130/85mmHg	35	38
	yes	32	34
Blood pressure	No	18	16
	below 130/85mmHg	15	12
TYPE II DIABETES	above 130/85mmHg	35	38
	yes	32	34
	No	18	16




Fig. 3. Significant changes in parameters in statin therapy




Fig. 4 . Comparison of Mets in both group A and group B.

Table 4 . Comparison of Mets in both group A and group B

Source	Atorvastatin (no of patients)	Rosuvastatin (no of patients)
Mets established	33 (66%)	26 (52%)
Mets not established	17 (34%)	24 (48%)

Group B, 23 were male, and 27 were female. Females are affected more than males. Data from two groups were analysed for an efficacy comparison in the intention-to-treat population. Baseline values of all parameters were similar between the two groups. The mean Serum TC, TG, LDL-C, and VLDL-C levels were significantly reduced on therapy.

4. Discussion

Metabolic syndrome is a disorder that increases the risk of CHD [10-14]. During 6 months period, a total of 100 participants were collected. Collected patients were divided into two groups based on their treatment plan as Group A (Atorvastatin) and Group B (Rosuvastatin). More participants were observed in the age group above 60 yrs. Among 100 patients, out of 50 patients belonging to Group A, 24 were male, and 26 were female. Out of 50 patients belonging to

Coming to the lipid profile, in group A, the mean triglyceride levels before and after treatments were found to be 214.2±0.86 and 185.94±1.22, respectively. Mean HDL levels before and after treatment were found to be 31.52±0.35, and 39.92±0.46 respectively. Mean LDL values before treatment are 161.4±1.09 and after treatment is 144.96±0.68. Whereas the mean VLDL values before and after treatment are 35.98±0.56 and 27.14±0.21 respectively.

In group B, the mean triglyceride levels before and after treatments were found to be 215.98 ± 0.62 and 142.04 ± 0.30 respectively. Mean HDL levels before and after treatment were found to be 28.34 ± 0.48 and 42.04 ± 0.30 respectively. Mean LDL values before treatment are 163.16 ± 0.94 and after treatment is 138.34 ± 0.73 . Whereas the mean VLDL values before and after treatment are 36.12 ± 0.43 and 24.72 ± 0.27 respectively.

Rosuvastatin reduced the level of TC by 215.98 ± 0.62 to 42.04 ± 0.30 ($p < 0.0001$) and LDL by 163.16 ± 0.94 to 138.34 ± 0.73 ($p < 0.01$) and VLDL-C by 36.12 ± 0.43 to 24.72 ± 0.27 ($p = 0.0001$). The level of HDL was from 28.34 ± 0.48 to 42.04 ± 0.30 ($p = 0.0001$). After 3 months of treatment, greater reductions in total cholesterol (181.48 ± 1.98 vs 174.32 ± 2.08), LDL (144.96 ± 0.68 vs 138.34 ± 0.73), triglycerides (185.94 ± 1.22 vs 181.74 ± 1.77), and increased high-density lipoprotein (39.92 ± 0.46 vs 42.04 ± 0.3) levels were observed in the atorvastatin group and Rosuvastatin group respectively. The Rosuvastatin group ($P = 0.01$) exhibited a significantly greater reduction as compared to the atorvastatin group ($P = 0.002$) [15, 16].

Among patients with MetS, rosuvastatin therapy resulted in significantly higher low-density lipoprotein cholesterol and total cholesterol reduction as compared with Atorvastatin [17-20]. Similarly, the significantly higher percentage of patients receiving rosuvastatin therapy was successful in achieving the target of total cholesterol and triglycerides as compared to atorvastatin [21, 22]. This resulted in reducing the risk of metabolic syndrome in patients receiving rosuvastatin (48%) than atorvastatin (34%) [17, 23].

5. Conclusion

In the present study, both Atorvastatin and Rosuvastatin were well tolerated in lowering cholesterol. This study revealed a significant reduction in lipid levels (total cholesterol, and LDL), an increase in HDL levels was achieved by rosuvastatin and atorvastatin respectively. An intermediate significant difference in the reduction of other lipid levels was observed between Atorvastatin and Rosuvastatin after 3 months of treatment. Metabolic syndrome is

a constellation of cardiac risk factors that increases CVD risk. Statins have been repeatedly shown to decrease risk in a variety of patient groups. There are no studies directly addressing statin treatment in metabolic syndrome, but the present study confirmed that rosuvastatin therapy is commonly prescribed doses is the most effective statin for low-density lipoprotein cholesterol goal achievement and for improving the lipid profile in with MetS.

Conflict of Interest

The authors hereby declare that they have no conflict of interest.

Author's contributions

All authors equally participated in designing experiment analysis and interpretation of data. All authors read and approved the final manuscript.

Consent for publications

All authors have read and approved the final manuscript for publication.

Availability of data and material

The authors have embedded all data in the manuscript.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Ethics approval and consent to participate

The authors did not use human or animals in the research

References

1. Niwa K (2021) Metabolic syndrome and coronary artery disease in adults with congenital heart disease. Cardiovascular diagnosis and therapy 11 (2): 563-576. doi:<https://doi.org/10.21037/cdt-20-781>
2. Lin T-Y, Chien K-L, Chiu Y-H, Chuang P-C, Yen M-F, Chen H-H (2021) Dynamics of detailed components of metabolic syndrome associated with the risk of cardiovascular disease and death. Scientific Reports 11 (1): 3677.

- doi:<https://doi.org/10.1038/s41598-021-83118-y>
3. Bhalwar R (2020) Metabolic syndrome: The Indian public health perspective. *Medical Journal Armed Forces India* 76 (1): 8-16. doi:<https://doi.org/10.1016/j.mjafi.2019.12.001>
 4. Yoon SJ, Kim SK, Lee NY, Choi YR, Kim HS, Gupta H, Youn GS, Sung H, Shin MJ, Suk KT (2021) Effect of Korean Red Ginseng on metabolic syndrome. *Journal of Ginseng Research* 45 (3): 380-389. doi:<https://doi.org/10.1016/j.jgr.2020.11.002>
 5. Na Z, Jiang H, Meng Y, Song J, Feng D, Fang Y, Shi B, Li D (2022) Association of galactose and insulin resistance in polycystic ovary syndrome: A case-control study. *eClinicalMedicine* 47): 101379. doi:<https://doi.org/10.1016/j.eclinm.2022.101379>
 6. Herrett E, Strongman H, Gadd S, Tomlinson L, Nitsch D, Bhaskaran K, Williamson E, van Staa T, Sofat R, Timmis A, Wells S, Smeeth L, Jackson R (2022) The importance of blood pressure thresholds versus predicted cardiovascular risk on subsequent rates of cardiovascular disease: a cohort study in English primary care. *The Lancet Healthy Longevity* 3 (1): e22-e30. doi:[https://doi.org/10.1016/S2666-7568\(21\)00281-6](https://doi.org/10.1016/S2666-7568(21)00281-6)
 7. Handelsman Y, Jellinger PS, Guerin CK, Bloomgarden ZT, Brinton EA, Budoff MJ, Davidson MH, Einhorn D, Fazio S, Fonseca VA, Garber AJ, Grunberger G, Krauss RM, Mechanick JL, Rosenblit PD, Smith DA, Wyne KL (2020) Consensus Statement by the American Association of Clinical Endocrinologists and American College of Endocrinology on the Management of Dyslipidemia and Prevention of Cardiovascular Disease Algorithm – 2020 Executive Summary. *Endocrine Practice* 26 (10): 1196-1224. doi:<https://doi.org/10.4158/CS-2020-0490>
 8. Alberti KGMM, Zimmet PZ (1998) Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus. Provisional report of a WHO consultation. *Diabetic medicine* 15 (7): 539-553. doi:[https://doi.org/10.1002/\(SICI\)1096-9136\(199807\)15:7%3C539::AID-DIA668%3E3.0.CO;2-S](https://doi.org/10.1002/(SICI)1096-9136(199807)15:7%3C539::AID-DIA668%3E3.0.CO;2-S)
 9. Music M, Dervisevic A, Pepic E, Lepara O, Fajkic A, Ascic-Buturovic B, Tuna E (2015) Metabolic Syndrome and Serum Liver Enzymes Level at Patients with Type 2 Diabetes Mellitus. *Medical archives (Sarajevo, Bosnia and Herzegovina)* 69 (4): 251-255. doi:<https://doi.org/10.5455/medarh.2015.69.251-255>
 10. Kachur S, Morera R, De Schutter A, Lavie CJ (2018) Cardiovascular Risk in Patients with Prehypertension and the Metabolic Syndrome. *Current Hypertension Reports* 20 (2): 15. doi:<https://doi.org/10.1007/s11906-018-0801-2>
 11. Ansarimoghaddam A, Adineh HA, Zareban I, Iranpour S, HosseinZadeh A, Kh F (2018) Prevalence of metabolic syndrome in Middle-East countries: Meta-analysis of cross-sectional studies. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* 12 (2): 195-201. doi:<https://doi.org/10.1016/j.dsx.2017.11.004>
 12. Zhao Y, Evans MA, Allison MA, Bertoni AG, Budoff MJ, Criqui MH, Malik S, Ouyang P, Polak JF, Wong ND (2019) Multisite atherosclerosis in subjects with metabolic syndrome and diabetes and relation to cardiovascular events: The Multi-Ethnic Study of Atherosclerosis. *Atherosclerosis* 282): 202-209. doi:<https://doi.org/10.1016/j.atherosclerosis.2018.12.005>
 13. Haswell C, Ali A, Page R, Hurst R, Rutherford-Markwick K (2021) Potential of Beetroot and Blackcurrant Compounds to Improve Metabolic Syndrome Risk Factors. *Metabolites* 11 (6): 338. doi:<https://doi.org/10.3390/metabo11060338>
 14. Alshammary AF, Alharbi KK, Alshehri NJ, Vennu V, Ali Khan I (2021) Metabolic syndrome and coronary artery disease risk: a meta-analysis of observational studies. *International Journal of Environmental Research and Public Health* 18 (4): 1773. doi:<https://doi.org/10.3390/ijerph18041773>

15. Werida R, Khairat I, Khedr NF (2021) Effect of atorvastatin versus rosuvastatin on inflammatory biomarkers and LV function in type 2 diabetic patients with dyslipidemia. *Biomedicine & Pharmacotherapy* 135): 111179. doi:<https://doi.org/10.1016/j.biopha.2020.111179>
16. Kummen M, Solberg OG, Storm-Larsen C, Holm K, Ragnarsson A, Trøseid M, Vestad B, Skårdal R, Yndestad A, Ueland T, Svardal A, Berge RK, Seljeflot I, Gullestad L, Karlsen TH, Aaberge L, Aukrust P, Hov JR (2020) Rosuvastatin alters the genetic composition of the human gut microbiome. *Scientific Reports* 10 (1): 5397. doi:<https://doi.org/10.1038/s41598-020-62261-y>
17. Fahmy MIM, Sayed RH, El-Yamany MF, El-Naggar R, A. Eliwa H (2022) Rosuvastatin and co-enzyme Q10 improve high-fat and high-fructose diet-induced metabolic syndrome in rats via ameliorating inflammatory and oxidative burden. *Biomedicine & Pharmacotherapy* 153): 113526. doi:<https://doi.org/10.1016/j.biopha.2022.113526>
18. Al-Sabti H, Al-Hinai AT, Al-Zakwani I, Al-Hashmi K, Al Mahmeed W, Arafah M, Shehab A, Al Tamimi O, Al Awadhi M, Al Herz S, Al Anazi F, Al Nemer K, Metwally O, Alkhadra A, Fakhry M, Elghetany H, Medani AR, Yusufali AH, Al Jassim O, Al Hallaq O, Baslaib F, Amin H, Al-Waili K, Al-Rasadi K (2022) The Achievement of Non-high-density Lipoprotein Cholesterol Target in Patients with Very High Atherosclerotic Cardiovascular Disease Risk Stratified by Triglyceride Levels Despite Statin-controlled Low-density Lipoprotein Cholesterol. *Oman medical journal* 37 (2): e367. doi:<https://doi.org/10.5001/omj.2022.79>
19. Gu A, Kamat S, Argulian E (2018) Trends and disparities in statin use and low-density lipoprotein cholesterol levels among US patients with diabetes, 1999–2014. *Diabetes Research and Clinical Practice* 139): 1-10. doi:<https://doi.org/10.1016/j.diabres.2018.02.019>
20. Santos HO, Earnest CP, Tinsley GM, Izidoro LFM, Macedo RCO (2020) Small dense low-density lipoprotein-cholesterol (sdLDL-C): Analysis, effects on cardiovascular endpoints and dietary strategies. *Progress in Cardiovascular Diseases* 63 (4): 503-509. doi:<https://doi.org/10.1016/j.pcad.2020.04.009>
21. Chilbert MR, VanDuyn D, Salah S, Clark CM, Ma Q (2022) Combination Therapy of Ezetimibe and Rosuvastatin for Dyslipidemia: Current Insights. *Drug Des Devel Ther* 16): 2177-2186. doi:<https://doi.org/10.2147/dddt.s332352>
22. Rizos CV, Skoumas I, Rallidis L, Skolidis E, Tziomalos K, Garoufi A, Anagnostis P, Sfikas G, Kotsis V, Doumas M, Kolovou G, Lambadiari V, Dima I, Kiouri E, Zacharis E, Agapakis D, Attilakos A, Antza C, Vlachopoulos C, Liberopoulos EN (2021) LDL cholesterol target achievement in heterozygous familial hypercholesterolemia patients according to 2019 ESC/EAS lipid guidelines: Implications for newer lipid-lowering treatments. *International Journal of Cardiology* 345): 119-124. doi:<https://doi.org/10.1016/j.ijcard.2021.10.024>
23. Tabrizi R, Tamtaji OR, Mirhosseini N, Lankarani KB, Akbari M, Dadgostar E, Borhani-Haghighi A, Peymani P, Ahmadizar F, Asemi Z (2019) The effects of statin use on inflammatory markers among patients with metabolic syndrome and related disorders: A systematic review and meta-analysis of randomized controlled trials. *Pharmacological Research* 141): 85-103. doi:<https://doi.org/10.1016/j.phrs.2018.12.010>



Copyright © 2023 by the author(s). This is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

How to Cite This Article:

Raju KR, Sharifa A, Soumya P, Khanam R, Bhargav KkBS (2023) Comparative study on the efficacy of statin therapy in lowering cholesterol and assessing the risk of metabolic syndrome in patients under statins therapy. Cellular, Molecular and Biomedical Reports 3 (1): 1-8. doi:10.55705/cmbr.2022.361364.1064

Download citation:

[RIS](#); [EndNote](#); [Mendeley](#); [BibTeX](#); [APA](#); [MLA](#); [HARVARD](#); [VANCOUVER](#)